

Date: 18.09.2021; 1400.06.27

Patient's Name: F.Gh

Responsible Physician: Dr.Omranipour

Patient presentation:

A 51 year-old woman with right breast locally advanced cancer who underwent neoadjuvant chemotherapy (8 sessions)

PEx: locally advanced (T4) and right axillary ulcerated fix mass

MG 99.7: suspicious spiculated right UOQ mass (B5) + suspicious LAP

US: Right breast mass (B5)

CT: Right breast mass (70mm) + Right axillary mass (45mm), No metastasis

Bone Scan: 6/7th ribs are suspicious

Breast CNB: IDC, G2, ER+ (10-20%), PR+ (20-30%), HER2 -, Ki67 30%

LN FNA: positive

MRV: Right axillary vein thrombosis, pectoralis major muscle involvement and right retro hemi thoracic mass




Question:

What's the next plan?

Systemic or local treatment?

Recommendation:

1. Left T7 involvement & thoracic mass with LD involvement (in imaging & MRV review)
2. She is not operable
3. Radiotherapy is not helpful because of ROM limitation
4. She should be referred to medical oncologist (continue systemic therapy with some other medication)



پستون
مركز
بيماري

Research Centre

مرکز تحقیقات بیماریهای پستان



Breast Disease Research Center